DEPAR		ER UNIVERSITY EMATICS AND STATISTICS			
	Final	Exam			
	MAT	H 232			
December 15, 1999, 3:30-6:30 p.m.					
Name:					
	family name	given name			
Number:					

#### INSTRUCTIONS

- 1. This exam has 14 questions on 16 pages. Please check to make sure your exam is complete.
- 2. Write your final answer in the answer box.
- In each question indicate how you obtain your answer.
   You may lose points if your work is poorly presented.
- 4. If you need more room, use the reverse side of the **previous page** to show your work.
- 5. No calculators or other computing devices may be used.
- 6. Please write with a black or blue pen.

Question	Score	Max
1		6
2		6
3		6
4		7
5		5
6		6
7		10
8		8
9		4
10		11
11		9
12		8
13		7
14		7
Total		100

[4] 1. (a) Give a precise description of the kinds of row operation which are permitted in bringing a matrix to reduced row-echelon form.

ANSWER		

[2] (b) **Define an elementary matrix**.

ANSWER

ROUGH WORK IF REQUIRED

[6] 2	2. Find a basis for the set of solutions to the system	ANSWER
	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	

3. Let  $V = \mathsf{sp}(\boldsymbol{a}_1, \boldsymbol{a}_2, \boldsymbol{a}_3, \boldsymbol{a}_4, \boldsymbol{a}_5)$  denote the subspace of  $\mathbb{R}^5$  spanned by

$$a_{1} = [2, 3, 1, -1, 0]$$

$$a_{2} = [-1, 1, 1, 2, 2]$$

$$a_{3} = [7, 3, -1, -8, -6]$$

$$a_{4} = [0, 10, 6, 6, 8]$$

$$a_{5} = [1, -1, 1, -1, -3]$$

and  $A = \begin{bmatrix} a_1 & a_2 & a_3 & a_4 & a_5 \end{bmatrix}$  be the  $5 \times 5$  matrix whose columns are  $a_1, a_2, a_3, a_4, a_5$ .

By elementary row operations  $\boldsymbol{A}$  is converted to

	1	0	$\begin{array}{c} 2\\ -3\\ 0\\ 0\\ 0\\ 0\end{array}$	2	0	1
	0	1	-3	4	0	
H =	0	0	0	0	1	
	0	0	0	0	0	
	0	0	0	0	0	

[2]	(a)	Write down a	basis for	V.
-----	-----	--------------	-----------	----

[2] (b) Write down a basis for the row space of A.

[2] (c) Determine the rank of *A*. Give a reason for your answer.



## ROUGH WORK IF REQUIRED

[7] 4. On a separate sheet circulated with this exam you find the definition of a vector space over  $\mathbb R$  .

Let V be a vector space over  $\mathbb{R}$ . From the axioms listed in the definition, prove that, for any two vectors v and w in V there exists a unique vector x in V such that v + x = w.

ANSWER

ROUGH WORK

[5] 5. Let V be a vector space over  $\mathbb{R}$ . Let  $W_1$  and  $W_2$  be two subspaces of V. Prove that their intersection  $W_1 \cap W_2$  is a subspace of V.

ANSWER	

ROUGH WORK

[6] 6. Let  $\mathbb{R}^{2 \times 2}$  denote the vector space of all  $2 \times 2$  real matrices, using as vector addition and scalar multiplication the usual addition of matrices and multiplication of a matrix by a scalar.

Given are four matrices

$$\boldsymbol{v}_1 = \begin{bmatrix} 1 & 2 \\ -1 & 2 \end{bmatrix}, \quad \boldsymbol{v}_2 = \begin{bmatrix} 0 & 2 \\ -1 & 4 \end{bmatrix}, \quad \boldsymbol{v}_3 = \begin{bmatrix} 0 & -1 \\ 3 & 1 \end{bmatrix}, \quad \boldsymbol{v}_4 = \begin{bmatrix} 0 & -3 \\ 2 & 0 \end{bmatrix}.$$

It is given that  $\mathcal{B}=(m{v}_1,\ m{v}_2,\ m{v}_3,\ m{v}_4)$  is an ordered basis for  $\mathbb{R}^{2 imes 2}$ 

Let

$$v = \left[ egin{array}{cc} 2 & -1 \ 8 & 3 \end{array} 
ight]$$

Find the coordinate vector  $v_{\mathcal{B}}$  of v relative to  $\mathcal{B}$ .

ANSWER

7. Let F be the vector space of all functions mapping  $\mathbb{R}$  to  $\mathbb{R}$ . Let W be the subspace of F spanned by the four functions 1, x,  $e^x$  and  $xe^x$ . It is given that  $\mathcal{B} = (1, x, e^x, xe^x)$  is an ordered basis for W.

Given are two linear transformations  $T_1~:~W\to W$  and  $T_2~:~W\to W$  defined by

 $T_1(f) = f'$  (the derivative of f with respect to x) for all  $f \in W$ 

 $T_2(f) = f''$  (the second derivative of f with respect to x) for all  $f \in W$ .

Let  $A_1$  be the matrix representation of  $T_1$  relative to  $\mathcal{B}, \mathcal{B}$  and let  $A_2$  be the matrix representation of  $T_2$  relative to  $\mathcal{B}, \mathcal{B}$ .

[5]

(a) Find the matrix  $A_1$ . ANSWER

(Question 7. continues here.)

[3] (b) Decide whether the transformation  $T_1$  is invertible. Justify your answer.

ANSWER			

[2] (c) Use the composition of linear transformations to discover a simple relation between  $A_2$  and  $A_1$ . Justify your answer. Do not compute  $A_2$  explicitly, just express it in terms of  $A_1$ .

ANSWER

ROUGH WORK IF REQUIRED

ANSWER

[4] 8. (a) Given are three points P = (3, -1), Q = (2, 2) and R = (-1, 5). Find the area of the triangle PQR.

#### [4] (b) State the row-interchange property for determinants of square matrices. Use it to prove:

If two rows of a square matrix A are equal, then det(A) = 0.

ANSWER

# [4] 9. Evaluate the determinant

$$\begin{vmatrix} 0 & 2 & 3 & 5 \\ 1 & 0 & -2 & 1 \\ 0 & 0 & 3 & -4 \\ -3 & 0 & 1 & 7 \end{vmatrix}$$

ANSWER

10. Let

$$A = \left[ \begin{array}{rrr} 1 & -2 & 1 \\ 0 & 1 & -1 \\ 0 & 2 & -2 \end{array} \right] \; .$$

[5] (a) Find the eigenvalues and corresponding eigenspaces of A.



(Question 10. continues here.)


- [3]
- (b) Use diagonalization to compute  $A^{2000}$ . Give your answer in the form of a single  $3 \times 3$ matrix.

to ANSWER ive the × 3

[3] (c) Decide whether the matrix

$$B = \left[ \begin{array}{cc} 3 & -2 \\ 0 & 3 \end{array} \right]$$

is diagonalizable. Justify your answer.



[2] **11.** (a) Let a = [1, 2, -1] and b = [-1, 3, 0]. Find the projection of b on sp(a).

[3] (b) Let W be the subspace of  $\mathbb{R}^3$  defined by  $W = \{ [x, y, z] \in \mathbb{R}^3 \mid x - y + z = 0 \}.$ 

Write down the basis for  $W^{\perp},$  the orthogonal complement of W.

[4]

(c) Let c = [2, 1, 5]. Find the projection of c on W.

SHOW YOUR WORK (use the back of the previous page if necessary)

ANSWER

ANSWER	

**12.** Let  $V = sp(a_1, a_2, a_3)$  be the subspace of  $\mathbb{R}^4$  spanned by the vectors  $a_1 = [1, 0, 0, 1]$ ,  $a_2 = [1, 1, 0, 1]$  and  $a_3 = [0, 1, -1, 0]$ .

[6]	(a)	Find an orthogonal basis for $V$ .	ANSWER
[2]	(b)	Use your answer to part (a) to find an or- thonormal basis for V.	ANSWER

SHOW YOUR WORK (use the back of the previous page if necessary)

[7] 13. The following data points are given:

(-1,0), (0,1), (1,3), (2,6).

Find the least-squares linear fit for these data points.

ANSWER

14. Let  $E = (e_1, e_2)$  be the standard ordered basis for  $\mathbb{R}^2$ . Let  $b_1 = [2, -1]$ ,  $b_2 = [-3, 2]$  and let  $B = (b_1, b_2)$  be an ordered basis for  $\mathbb{R}^2$ .

Let  $T : \mathbb{R}^2 \to \mathbb{R}^2$  be the linear transformation defined by  $T([x_1, x_2]) = [x_1 + x_2, -x_1 + x_2]$  for every  $[x_1, x_2] \in \mathbb{R}^2$ .

### [2] (a) Write down the standard matrix representation of T.

[2]	(b)	Find the change-of-coordinates matrix from $E$ to $B$ .

SHOW YOUR WORK

ANSWER

ANSWER