

Three-phase Golay sequence and array triads

Aki Ayukawa Avis

Jonathan Jedwab

12 October 2019

Abstract

3-phase Golay sequence and array triads are a natural generalisation of 2-phase Golay sequence pairs, yet their study has until now been largely neglected. We present exhaustive search results for 3-phase Golay sequence triads for all lengths up to 24, showing that the existence pattern is much richer than that for 2-phase Golay sequence pairs. We give an elementary proof that there is no 3-phase Golay sequence triad whose length is congruent to 4 modulo 6. We give two construction methods for 3-phase Golay array triads, and show how to project 3-phase Golay array triads to lower-dimensional Golay triads. In this way we explain much of the existence pattern found by exhaustive search.

1 Introduction

Golay sequence pairs were introduced by Golay in 1951 to solve a problem in multislit spectrometry [11] and have since been applied in various digital communications schemes, including coded aperture imaging [17], optical time domain reflectometry [15], power control for multicarrier wireless transmission [5], and medical ultrasound [16]. They are related to Barker sequences [13] and Reed-Muller codes [5].

We consider a *length s sequence* to be an 1-dimensional matrix $\mathcal{A} = (A_i)$ of complex-valued entries, indexed by an integer i , for which

$$A_i = 0 \text{ if either } i < 0 \text{ or } i \geq s.$$

J. Jedwab is with Department of Mathematics, Simon Fraser University, 8888 University Drive, Burnaby BC V5A 1S6, Canada.

A. Avis is with Department of Nutrition, Université de Montréal, 2900 Boulevard Édouard-Montpetit, Montréal QB H3T 1J4, Canada.

Both authors were supported by NSERC.

Email: jed@sfu.ca, aki.avis@umontreal.ca

The *aperiodic autocorrelation function* of $\mathcal{A} = (A_i)$ is

$$C_{\mathcal{A}}(u) = \sum_i A_i \overline{A_{i+u}} \quad \text{for integers } u,$$

where bar represents complex conjugation. A 2-phase Golay sequence pair comprises length s sequences $\mathcal{A} = (A_i)$ and $\mathcal{B} = (B_i)$ whose elements lie in the alphabet $\{1, -1\}$, satisfying

$$(C_{\mathcal{A}} + C_{\mathcal{B}})(u) = 0 \quad \text{for all } u \neq 0. \quad (1.1)$$

Golay sequence pairs have been constructed for all lengths s of the form $2^a 10^b 26^c$, where a, b, c are non-negative integers [21], but no examples are known for other lengths. In view of the wide range of potential applications for such sequences, the definition of Golay sequence pairs has been extended in various ways: from the 2-phase alphabet $\{1, -1\}$ to the H -phase alphabet (comprising the H^{th} roots of unity) for even H [4, 5, 8, 18], and to quadrature amplitude modulation constellations [3, 19]; from sequence pairs to sequence sets of even size greater than 2 [18, 20]; and from sequences to arrays of dimension 2 or greater [6, 7, 14, 17].

Nonetheless, there is a natural generalisation of 2-phase Golay sequence pairs that, with the exception of a brief study by Frank [9] in 1980, appears to have been largely neglected: from the 2-phase alphabet $\{1, -1\}$ to the 3-phase alphabet $\{1, \omega, \omega^2\}$ where $\omega = e^{2\pi i/3}$, and simultaneously from sequence pairs to sequence triads (sets of size 3). The 3-phase alphabet is attractive for applications, because the small number of phases allows the received signal levels to be easily distinguished at the receiver. The related question of the existence of 3-phase Barker arrays was considered in [2].

We shall consider the following questions theoretically and computationally, summarising and extending results contained in the 2008 Master's thesis of the first author [1]:

1. For which lengths s do 3-phase Golay sequence triads exist?
2. How many 3-phase length s Golay sequence triads are there?
3. How are 3-phase Golay array triads related to 3-phase Golay sequence triads?
4. How can 3-phase Golay sequence and array triads be constructed?

Frank [9] gave an example of a 3-phase Golay sequence triad of length 3 and one of length 5, and presented a construction for a 3-phase length $3s$ Golay sequence triad from one of length s . Some of the other constructions described in [9] can be applied to 3-phase Golay sequence triads, but in general they produce sequences over larger alphabets such as 6-phase or 12-phase.

We shall show in Theorem 10 that the projection to lower dimensions of a multi-dimensional 3-phase Golay array triad is also a 3-phase Golay triad, so that a higher-dimensional Golay array triad can be considered a more fundamental object than its lower-dimensional projections; this parallels the central argument of [7] for Golay pairs. Our

main construction method for Golay array triads is Theorem 13, which modifies Frank's construction [9] for Golay sequence triads to produce an $(r + 1)$ -dimensional Golay array triad from an r -dimensional Golay array triad.

The rest of this paper is organised as follows. In Section 2, we establish structural properties of 3-phase Golay sequence triads, and in Table 1 present counts of 3-phase length s Golay sequence triads obtained from exhaustive search for each $s \leq 24$. In Section 3, motivated by computational evidence, we give an elementary proof that there is no 3-phase length s Golay sequence triad when $s \equiv 4 \pmod{6}$. In Section 4, we introduce 3-phase Golay array triads and show how they can be projected to 3-phase Golay array triads in one fewer dimension, deducing counts of 3-phase Golay array triads in Table 2 from those for sequence triads given in Table 1. In Section 5, we present two constructions of 3-phase Golay array triads, and apply them to a small set of seed triads to explain the existence of many of the triads counted in Tables 1 and 2. In Section 6, we pose some open questions motivated by our results. In the Appendix, we explicitly list some of the Golay sequence and array triads whose existence we were not able to explain.

2 Golay sequence triads

In this section, we establish structural properties of 3-phase Golay sequence triads and present exhaustive search results for each length up to 24.

Throughout, let $\omega = e^{2\pi i/3}$. A length s sequence (A_i) is a *3-phase sequence* if all elements of the set $\{A_i \mid 0 \leq i < s\}$ take values in $\{1, \omega, \omega^2\}$. An unordered set of three 3-phase length s sequences $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a *Golay sequence triad* if

$$(C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u) = 0 \quad \text{for all } u \neq 0. \quad (2.1)$$

It is sufficient to verify (2.1) for the integers u satisfying $0 < u < s$, because every complex-valued sequence \mathcal{A} satisfies $C_{\mathcal{A}}(-u) = \overline{C_{\mathcal{A}}(u)}$ for all integers u .

A 3-phase length s sequence (A_i) corresponds to a length s sequence (a_i) over \mathbb{Z}_3 , where

$$A_i = \omega^{a_i} \text{ for each } i \text{ satisfying } 0 \leq i < s.$$

The aperiodic autocorrelation function of a sequence over \mathbb{Z}_3 is that of the corresponding 3-phase sequence.

Example 1. *The sequences*

$$\begin{aligned} \mathcal{A} = (A_i) &= [1 \quad \omega^2 \quad 1 \quad 1 \quad \omega^2 \quad 1], \\ \mathcal{B} = (B_i) &= [1 \quad \omega \quad \omega^2 \quad \omega^2 \quad \omega^2 \quad \omega], \\ \mathcal{C} = (C_i) &= [1 \quad \omega \quad \omega \quad \omega \quad 1 \quad \omega^2], \end{aligned}$$

satisfy

$$\begin{aligned} (C_{\mathcal{A}}(u) \mid 0 \leq u < 6) &= [6 \quad -1 \quad 1 \quad 3 \quad -1 \quad 1], \\ (C_{\mathcal{B}}(u) \mid 0 \leq u < 6) &= [6 \quad -\omega \quad \omega \quad \omega - 1 \quad -\omega^2 \quad \omega^2], \\ (C_{\mathcal{C}}(u) \mid 0 \leq u < 6) &= [6 \quad -\omega^2 \quad \omega^2 \quad \omega^2 - 1 \quad -\omega \quad \omega], \end{aligned}$$

and therefore comprise a 3-phase length 6 Golay sequence triad. The corresponding sequences over \mathbb{Z}_3 are $(a_i) = [0 \ 2 \ 0 \ 0 \ 2 \ 0]$, $(b_i) = [0 \ 1 \ 2 \ 2 \ 2 \ 1]$, $(c_i) = [0 \ 1 \ 1 \ 1 \ 0 \ 2]$.

We will use upper-case letters for 3-phase sequences (“multiplicative notation”), and lower-case letters for sequences over \mathbb{Z}_3 (“additive notation”); the same letter (for example A and a) will indicate corresponding sequences. We will switch between multiplicative and additive notation as convenient.

Suppose that $\{(a_i), (b_i), (c_i)\}$ is a length s Golay sequence triad over \mathbb{Z}_3 . Since the sequences (x_i) and $(x_i + 1)$ over \mathbb{Z}_3 have identical aperiodic autocorrelation function, each of the 3^3 unordered sets $\{(a_i + \alpha), (b_i + \beta), (c_i + \gamma)\}$ is also a Golay sequence triad over \mathbb{Z}_3 , as α, β, γ range over $\{0, 1, 2\}$. We may therefore assume that

$$(a_0, b_0, c_0) = (0, 0, 0),$$

in which case the Golay sequence triad is in *normalised form*. Condition (2.1) with $u = s - 1$ then forces $\{a_{s-1}, b_{s-1}, c_{s-1}\} = \{0, 1, 2\}$, which implies there are exactly $3!$ ordered Golay sequence triads corresponding to each (unordered) Golay sequence triad.

The following result follows directly from the definition of aperiodic autocorrelation function and Golay sequence triad.

Lemma 2.

(i) (Linear Offsets). Suppose that $\{(a_i), (b_i), (c_i)\}$ is a length s Golay sequence triad over \mathbb{Z}_3 . Then

$$\{(a_i + ei), (b_i + ei), (c_i + ei)\}$$

is also a length s Golay sequence triad over \mathbb{Z}_3 for each $e \in \mathbb{Z}_3$.

(ii) (Reversal). Suppose that $\{(a_i), (b_i), (c_i)\}$ is a length s Golay sequence triad over \mathbb{Z}_3 . Then

$$\{(a_{s-1-i}), (b_{s-1-i}), (c_{s-1-i})\}$$

is also a length s Golay sequence triad over \mathbb{Z}_3 .

(iii) (Reverse Conjugation). Let (x_i) be a length s sequence over \mathbb{Z}_3 . Then the length s sequence $(2x_{s-1-i})$ over \mathbb{Z}_3 has identical aperiodic autocorrelation function to (x_i) .

Suppose that T is a length s Golay sequence triad over \mathbb{Z}_3 . By Lemma 2 (iii), when one sequence (x_i) of T is replaced by the length s sequence $(2x_{s-1-i})$, the resulting set is also a length s Golay sequence triad over \mathbb{Z}_3 . We say that the Golay sequence triads over \mathbb{Z}_3 obtained from T by applying one or more of the operations described in Lemma 2, and then taking the normalised form, are all *equivalent* to T . The size of the equivalence class of a normalised Golay sequence triad divides $3 \cdot 2 \cdot 2^3 = 48$. We take the representative of the equivalence class to be its lexicographically first member. For example, there are exactly three equivalence classes of length 5 Golay sequence triads over \mathbb{Z}_3 , each of size 24, and their equivalence class representatives are

$$\begin{aligned} & \{ [0 0 0 1 0], [0 1 2 2 1], [0 0 2 1 2] \}, \\ & \{ [0 0 1 0 0], [0 0 1 2 1], [0 0 2 1 2] \}, \\ & \{ [0 0 1 1 0], [0 0 1 2 1], [0 0 2 0 2] \}. \end{aligned}$$

Golay [12] showed that the elements of a 2-phase length s Golay sequence pair $\{(A_i), (B_i)\}$ are related by

$$A_i B_i = -A_{s-1-i} B_{s-1-i} \quad \text{for each } i \text{ satisfying } 0 \leq i < s.$$

We next derive a counterpart relationship for the elements of a 3-phase Golay sequence triad.

Proposition 3. *Suppose that $\{(A_i), (B_i), (C_i)\}$ is a 3-phase length s Golay sequence triad. Then*

$$A_i B_i C_i = A_{s-1-i} B_{s-1-i} C_{s-1-i} \quad \text{for each } i \text{ satisfying } 0 \leq i < s.$$

Proof. Let u satisfy $0 < u < s$. We are given that

$$\sum_{i=0}^{s-1-u} (A_i \overline{A_{i+u}} + B_i \overline{B_{i+u}} + C_i \overline{C_{i+u}}) = 0,$$

so the multiset of $3(s-u)$ summands above contains each of 1 , ω and ω^2 exactly $s-u$ times. The product of these summands is therefore

$$\prod_{i=0}^{s-1-u} (A_i \overline{A_{i+u}})(B_i \overline{B_{i+u}})(C_i \overline{C_{i+u}}) = 1^{s-u} \omega^{s-u} (\omega^2)^{s-u} = 1,$$

which implies that

$$\prod_{i=0}^{s-1-u} A_i B_i C_i = \prod_{i=u}^{s-1} A_i B_i C_i.$$

Since this holds for each u satisfying $0 < u < s$, the required result follows by induction on decreasing $u \leq s-1$. \square

The property of a length s Golay sequence triad $\{(a_i), (b_i), (c_i)\}$ over \mathbb{Z}_3 corresponding to Proposition 3 is

$$a_i + b_i + c_i \equiv a_{s-1-i} + b_{s-1-i} + c_{s-1-i} \pmod{3} \quad \text{for each } i \text{ satisfying } 0 \leq i < s. \quad (2.2)$$

We use this property to determine by exhaustive search the equivalence classes of length s Golay sequence triads $\{(a_i), (b_i), (c_i)\}$ over \mathbb{Z}_3 for each $s \leq 24$. We begin by fixing the outermost elements of the sequences of the triad: the normalisation condition gives $(a_0, b_0, c_0) = (0, 0, 0)$, and we may then assume from the autocorrelation condition $u = s-1$ of (2.1) that $(a_{s-1}, b_{s-1}, c_{s-1}) = (0, 1, 2)$. For each value of i in $\{1, 2, \dots, \lfloor \frac{s-1}{2} \rfloor\}$ taken in ascending order, we then recursively select all values of the outermost undetermined sequence elements $\{a_i, b_i, c_i, a_{s-1-i}, b_{s-1-i}, c_{s-1-i}\}$ that are consistent with both property (2.2) and the autocorrelation condition $u = s-1-i$ of (2.1). We finally collect the resulting Golay sequence triads into equivalence classes and retain only the representative of each equivalence class. (Since just the representative of each equivalence class is required, the search space can be further reduced: for example, by taking the leftmost nonzero entry of the subsequence (a_0, a_1, \dots, a_i) to be 1, and discarding search branches where this subsequence occurs lexicographically after the reverse conjugated subsequence $(2a_{s-1}, 2a_{s-2}, \dots, 2a_{s-1-i})$.) Table 1 shows the resulting counts of equivalence classes of Golay sequence triads, and of normalised Golay sequence triads. The existence pattern for 3-phase Golay sequence triads appears to be much richer than that for 2-phase Golay sequence pairs: within the range $1 \leq s \leq 24$, a 2-phase length s Golay sequence pair exists only for $s \in \{2, 4, 8, 10, 16, 20\}$ [21], whereas a 3-phase length s Golay sequence triad exists for all $s \notin \{4, 10, 16, 22\}$.

We call a sequence belonging to one or more 3-phase Golay sequence triads (not necessarily in normalised form) a *3-phase Golay sequence*. For example, each of the sequences $\mathcal{A}, \mathcal{B}, \mathcal{C}$ in Example 1 is a 3-phase length 6 Golay sequence. The first author discussed the use of 3-phase Golay sequences in an orthogonal frequency division multiplexing (OFDM) multicarrier transmission scheme, using their favourable peak-to-mean envelope power ratio properties to formulate an alternative exhaustive search algorithm for Golay sequence triads [1, Chap. 4] to the one described above. Since the transmission rate of an OFDM scheme using 3-phase Golay sequences increases with the number of Golay sequences of a given length, Table 1 also displays counts of Golay sequences.

3 Nonexistence result for $s \equiv 4 \pmod{6}$

A striking feature of Table 1 is that there are no 3-phase Golay sequence triads of length 4, 10, 16, and 22. In this section, we explain this observation by proving the following theorem.

Theorem 4. *There is no 3-phase length s Golay sequence triad when $s \equiv 4 \pmod{6}$.*

Sequence length	# equivalence classes						# normalised sequence triads	# Golay sequences
	size 1	size 8	size 16	size 24	size 48	Total		
2	1					1	1	9
3	1	1				2	9	27
4						0	0	0
5				3		3	72	108
6		3	3	4		10	168	288
7				8	9	17	624	792
8				1	3	4	168	306
9		15	33	25	32	105	2784	3708
10						0	0	0
11				14	50	64	2736	4932
12					7	7	336	756
13				16	48	64	2688	5850
14				23	68	91	3816	8334
15		36	306	51	126	519	12456	27576
16						0	0	0
17				10	15	25	960	2160
18		84	714	184	503	1485	40656	89532
19				6	11	17	672	1458
20					10	10	480	1080
21		84	2766	88	1007	3945	95376	202932
22						0	0	0
23				2		2	48	108
24		12	750	16	247	1025	24336	54756

Table 1: Counts of 3-phase length s Golay sequence triads, for each $s \leq 24$

Let $\mathcal{A} = (A_i)$ be a length s sequence. The *periodic autocorrelation function* of \mathcal{A} is

$$R_{\mathcal{A}}(u) = \sum_{i=0}^{s-1} A_i \overline{A_{(i+u) \bmod s}} \quad \text{for } 0 \leq u < s,$$

so that

$$R_{\mathcal{A}}(u) = C_{\mathcal{A}}(u) + \overline{C_{\mathcal{A}}(s-u)} \quad \text{for } 0 < u < s \quad (3.1)$$

by the definition of aperiodic autocorrelation. An unordered set of three 3-phase length s sequences $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a *periodic Golay sequence triad* if

$$(R_{\mathcal{A}} + R_{\mathcal{B}} + R_{\mathcal{C}})(u) = 0 \quad \text{for all } u \text{ satisfying } 0 < u < s.$$

From (3.1), the sequences of a Golay sequence triad also form a periodic Golay sequence triad of the same length.

Now we cannot prove Theorem 4 by attempting to rule out the existence of a 3-phase periodic Golay sequence triad of length congruent to 4 modulo 6: for example, the sequence set

$$\{ [1 \ 1 \ \omega \ \omega], [1 \ 1 \ \omega \ \omega], [1 \ \omega \ 1 \ \omega] \}.$$

is a periodic length 4 Golay sequence triad. However, we can constrain the elements of a 3-phase periodic Golay sequence triad as follows, which in combination with Proposition 3 will immediately prove Theorem 4.

Proposition 5. *Suppose that $\{(A_i), (B_i), (C_i)\}$ is a 3-phase length $2m$ periodic Golay sequence triad, where $m \equiv 2 \pmod{3}$. Then exactly two of the following three equations hold:*

$$\prod_{i=0}^{m-1} A_{2i} = \prod_{i=0}^{m-1} A_{2i+1}, \quad \prod_{i=0}^{m-1} B_{2i} = \prod_{i=0}^{m-1} B_{2i+1}, \quad \prod_{i=0}^{m-1} C_{2i} = \prod_{i=0}^{m-1} C_{2i+1}.$$

Proof. Let $\mathcal{A} = (A_i)$, $\mathcal{B} = (B_i)$, $\mathcal{C} = (C_i)$. We shall examine the value of $S \pmod{9}$, where

$$S = \left| \sum_{i=0}^{2m-1} (-1)^i A_i \right|^2 + \left| \sum_{i=0}^{2m-1} (-1)^i B_i \right|^2 + \left| \sum_{i=0}^{2m-1} (-1)^i C_i \right|^2. \quad (3.2)$$

For a length $2n$ sequence $\mathcal{X} = (X_i)$, we have

$$\begin{aligned} \left| \sum_{i=0}^{2n-1} (-1)^i X_i \right|^2 &= \sum_{i=0}^{2n-1} (-1)^i X_i \sum_{j=0}^{2n-1} (-1)^j \overline{X_j} \\ &= \sum_{i=0}^{2n-1} (-1)^i X_i \sum_{u=0}^{2n-1} (-1)^{i+u} \overline{X_{(i+u) \bmod 2n}} \end{aligned}$$

using the re-indexing $j = (i + u) \bmod 2n$, so that

$$\left| \sum_{i=0}^{2n-1} (-1)^i X_i \right|^2 = \sum_{u=0}^{2n-1} (-1)^u R_{\mathcal{X}}(u).$$

It follows from (3.2) that

$$\begin{aligned} S &= \sum_{u=0}^{2m-1} (-1)^u (R_{\mathcal{A}} + R_{\mathcal{B}} + R_{\mathcal{C}})(u) \\ &= (R_{\mathcal{A}} + R_{\mathcal{B}} + R_{\mathcal{C}})(0) \end{aligned}$$

because $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a periodic Golay triad. Since $(R_{\mathcal{A}} + R_{\mathcal{B}} + R_{\mathcal{C}})(0) = 6m$ and $m \equiv 2 \pmod{3}$, this implies that

$$S \equiv 3 \pmod{9}. \quad (3.3)$$

Now for a 3-phase length $2n$ sequence $\mathcal{X} = (X_i)$, by taking $Y_i = X_{2i}$ and $Z_i = X_{2i+1}$ in Lemma 6 given below, we find that $|\sum_{i=0}^{2n-1} (-1)^i X_i|^2$ is an integer satisfying the congruence

$$\left| \sum_{i=0}^{2n-1} (-1)^i X_i \right|^2 \equiv \begin{cases} 0 \pmod{9} & \text{if } \prod_{i=0}^{n-1} X_{2i} = \prod_{i=0}^{n-1} X_{2i+1}, \\ 3 \pmod{9} & \text{otherwise.} \end{cases}$$

The result then follows from (3.2) and (3.3). □

Lemma 6. *Let (Y_i) and (Z_i) be 3-phase length n sequences. Then $|\sum_{i=0}^{n-1} (Y_i - Z_i)|^2$ is an integer satisfying the congruence*

$$\left| \sum_{i=0}^{n-1} (Y_i - Z_i) \right|^2 \equiv \begin{cases} 0 \pmod{9} & \text{if } \prod_{i=0}^{n-1} Y_i = \prod_{i=0}^{n-1} Z_i, \\ 3 \pmod{9} & \text{otherwise.} \end{cases}$$

Proof. For $j = 0, 1, 2$, let the sequence (Y_i) contain ω^j exactly α_j times and let the sequence (Z_i) contain ω^j exactly β_j times. Then

$$\alpha_0 + \alpha_1 + \alpha_2 = \beta_0 + \beta_1 + \beta_2 = n, \quad (3.4)$$

and

$$\begin{aligned} \sum_{i=0}^{n-1} (Y_i - Z_i) &= (\alpha_0 - \beta_0) + \omega(\alpha_1 - \beta_1) + \omega^2(\alpha_2 - \beta_2) \\ &= a + \omega b, \end{aligned}$$

using $\omega^2 = -1 - \omega$ and writing $a = (\alpha_0 - \alpha_2) - (\beta_0 - \beta_2)$ and $b = (\alpha_1 - \alpha_2) - (\beta_1 - \beta_2)$. Therefore

$$\left| \sum_{i=0}^{n-1} (Y_i - Z_i) \right|^2 = a^2 + b^2 - ab,$$

which is an integer because a and b are integers. Since $a + b \equiv 0 \pmod{3}$ by (3.4), this implies

$$\left| \sum_{i=0}^{n-1} (Y_i - Z_i) \right|^2 \equiv 3b^2 \pmod{9}. \quad (3.5)$$

Now $\prod_{i=0}^{n-1} Y_i = 1^{\alpha_0} \omega^{\alpha_1} (\omega^2)^{\alpha_2} = \omega^{\alpha_1 - \alpha_2}$, and likewise $\prod_{i=0}^{n-1} Z_i = \omega^{\beta_1 - \beta_2}$, so that

$$\prod_{i=0}^{n-1} Y_i = \prod_{i=0}^{n-1} Z_i \quad \text{if and only if} \quad b \equiv 0 \pmod{3}$$

by the definition of b . Combination with (3.5) gives the required result. \square

Proof of Theorem 4. Suppose, for a contradiction, that $\{(A_i), (B_i), (C_i)\}$ is a 3-phase length $2m$ Golay sequence triad, where $m \equiv 2 \pmod{3}$. Then by Proposition 3,

$$\prod_{i=0}^{m-1} A_{2i} B_{2i} C_{2i} = \prod_{i=0}^{m-1} A_{2i+1} B_{2i+1} C_{2i+1}. \quad (3.6)$$

But $\{(A_i), (B_i), (C_i)\}$ is also a 3-phase length $2m$ periodic Golay sequence triad by (3.1), so we obtain a contradiction to (3.6) from Proposition 5. \square

4 Golay array triads

In this section, we extend the definition of a Golay sequence triad to multiple dimensions, and explore the relationship between multi-dimensional Golay array triads and Golay sequence triads.

We consider an $s_1 \times \cdots \times s_r$ array to be an r -dimensional matrix $\mathcal{A} = (A_{i_1, \dots, i_r})$ of complex-valued entries, indexed by integers i_1, \dots, i_r , for which

$$A_{i_1, \dots, i_r} = 0 \text{ if, for at least one } k \in \{1, \dots, r\}, \text{ either } i_k < 0 \text{ or } i_k \geq s_k.$$

The *aperiodic autocorrelation function* of \mathcal{A} is

$$C_{\mathcal{A}}(u_1, \dots, u_r) = \sum_{i_1, \dots, i_r} A_{i_1, \dots, i_r} \overline{A_{i_1+u_1, \dots, i_r+u_r}} \quad \text{for integers } u_1, \dots, u_r.$$

This function satisfies

$$C_{\mathcal{A}}(-u_1, \dots, -u_r) = \overline{C_{\mathcal{A}}(u_1, \dots, u_r)} \quad \text{for all } (u_1, \dots, u_r). \quad (4.1)$$

The following definitions are each analogous to those for sequences: a 3-phase array; the $s_1 \times \cdots \times s_r$ array (a_{i_1, \dots, i_r}) over \mathbb{Z}_3 corresponding to a 3-phase $s_1 \times \cdots \times s_r$ array (A_{i_1, \dots, i_r}) ; and the aperiodic autocorrelation function of (a_{i_1, \dots, i_r}) .

An unordered set of three 3-phase $s_1 \times \cdots \times s_r$ arrays $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a *Golay array triad* if

$$(C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1, \dots, u_r) = 0 \quad \text{for all } (u_1, \dots, u_r) \neq (0, \dots, 0),$$

and an array \mathcal{A} is called a *Golay array* if it belongs to one or more 3-phase Golay array triads.

Example 7. *The arrays*

$$\mathcal{A} = (a_{i,j}) = \begin{bmatrix} 0 & 0 & 2 \\ 2 & 0 & 0 \end{bmatrix}, \quad \mathcal{B} = (b_{i,j}) = \begin{bmatrix} 0 & 2 & 2 \\ 1 & 2 & 1 \end{bmatrix}, \quad \mathcal{C} = (c_{i,j}) = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 1 & 2 \end{bmatrix}$$

over \mathbb{Z}_3 satisfy

$$\begin{aligned} (C_{\mathcal{A}}(u, v) \mid 0 \leq u < 2, -3 < v < 3) &= \begin{bmatrix} -1 & 1 & 6 & 1 & -1 \\ 1 & -1 & 0 & 2 & 1 \end{bmatrix}, \\ (C_{\mathcal{B}}(u, v) \mid 0 \leq u < 2, -3 < v < 3) &= \begin{bmatrix} -\omega & \omega^2 & 6 & \omega & -\omega^2 \\ \omega & -\omega^2 & 0 & 2\omega & \omega^2 \end{bmatrix}, \\ (C_{\mathcal{C}}(u, v) \mid 0 \leq u < 2, -3 < v < 3) &= \begin{bmatrix} -\omega^2 & \omega & 6 & \omega^2 & -\omega \\ \omega^2 & -\omega & 0 & 2\omega^2 & \omega \end{bmatrix}, \end{aligned}$$

and therefore comprise a 2×3 Golay array triad over \mathbb{Z}_3 . The 2×3 array $(a_{i,j})$ can instead be represented as the 3×2 array $(a'_{i,j})$, where $a'_{i,j} = a_{j,i}$ for all i, j . However, we do not consider arrays obtained by reordering dimensions to be distinct: they are different formal representations of the same object.

Since the arrays (x_{i_1, \dots, i_r}) and $(x_{i_1, \dots, i_r} + 1)$ over \mathbb{Z}_3 have identical aperiodic autocorrelation function, an $s_1 \times \dots \times s_r$ Golay sequence triad $\{(a_{i_1, \dots, i_r}), (b_{i_1, \dots, i_r}), (c_{i_1, \dots, i_r})\}$ over \mathbb{Z}_3 may be assumed to satisfy

$$(a_{0, \dots, 0}, b_{0, \dots, 0}, c_{0, \dots, 0}) = (0, 0, 0),$$

in which case we say it is in *normalised form*.

The following result is the multi-dimensional version of Lemma 2.

Lemma 8.

(i) (Linear Offsets). *Suppose that $\{(a_{i_1, \dots, i_r}), (b_{i_1, \dots, i_r}), (c_{i_1, \dots, i_r})\}$ is an $s_1 \times \dots \times s_r$ Golay array triad over \mathbb{Z}_3 . Then*

$$\{(a_{i_1, \dots, i_r} + e_1 i_1 + \dots + e_r i_r), (b_{i_1, \dots, i_r} + e_1 i_1 + \dots + e_r i_r), (c_{i_1, \dots, i_r} + e_1 i_1 + \dots + e_r i_r)\}$$

is also an $s_1 \times \dots \times s_r$ Golay array triad over \mathbb{Z}_3 for all $e_1, \dots, e_r \in \mathbb{Z}_3$.

(ii) (Reversal). *Suppose that $\{(a_{i_1, \dots, i_r}), (b_{i_1, \dots, i_r}), (c_{i_1, \dots, i_r})\}$ is an $s_1 \times \dots \times s_r$ Golay array triad over \mathbb{Z}_3 . Then for each $k \in \{1, \dots, r\}$,*

$$\{(a_{i_1, \dots, s_k - 1 - i_k, \dots, i_r}), (b_{i_1, \dots, s_k - 1 - i_k, \dots, i_r}), (c_{i_1, \dots, s_k - 1 - i_k, \dots, i_r})\}$$

is also an $s_1 \times \dots \times s_r$ Golay array triad over \mathbb{Z}_3 .

(iii) (Reverse Conjugation). *Let (x_{i_1, \dots, i_r}) be an $s_1 \times \dots \times s_r$ array over \mathbb{Z}_3 . Then the $s_1 \times \dots \times s_r$ array $(2x_{s_1-1-i_1, \dots, s_r-1-i_r})$ over \mathbb{Z}_3 has identical aperiodic autocorrelation function to (x_{i_1, \dots, i_r}) .*

The Golay array triads over \mathbb{Z}_3 obtained by applying one or more of the operations described in Lemma 8, and then normalising, form an equivalence class whose size divides $3^r \cdot 2^r \cdot 2^3 = 2^{r+3} \cdot 3^r$. We take the representative of the equivalence class to be its lexicographically first member. For example, there are exactly three equivalence class of 2×7 Golay array triads over \mathbb{Z}_3 , each of size 288, and their equivalence class representatives are

$$\left\{ \left[\begin{array}{ccccccc} 0 & 0 & 0 & 1 & 0 & 2 & 2 \\ 0 & 0 & 2 & 0 & 1 & 1 & 0 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 2 & 0 & 1 & 2 & 0 & 0 \\ 0 & 2 & 2 & 2 & 0 & 2 & 1 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 1 & 0 & 2 & 1 & 1 & 1 \\ 0 & 1 & 2 & 2 & 2 & 0 & 2 \end{array} \right] \right\},$$

$$\left\{ \left[\begin{array}{ccccccc} 0 & 0 & 0 & 1 & 0 & 2 & 2 \\ 2 & 0 & 2 & 1 & 0 & 0 & 0 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 0 & 2 & 0 & 1 & 1 & 0 \\ 2 & 0 & 1 & 1 & 1 & 2 & 1 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 0 & 1 & 0 & 2 & 0 & 1 \\ 2 & 0 & 0 & 2 & 2 & 1 & 2 \end{array} \right] \right\},$$

$$\left\{ \left[\begin{array}{ccccccc} 0 & 0 & 0 & 1 & 0 & 2 & 2 \\ 2 & 2 & 0 & 2 & 1 & 2 & 0 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 1 & 1 & 1 & 1 & 0 & 2 \\ 1 & 1 & 2 & 0 & 1 & 2 & 1 \end{array} \right], \left[\begin{array}{ccccccc} 0 & 1 & 0 & 0 & 2 & 2 & 0 \\ 1 & 1 & 1 & 0 & 2 & 1 & 2 \end{array} \right] \right\}.$$

We now introduce an invertible mapping that reduces the number of dimensions of an array by exactly one and that maps Golay array triads to Golay array triads. Our method is modelled on that of [14] for 2-phase Golay array pairs, as subsequently used in [7] and [10] for H -phase Golay array pairs where H is even. For an $s_1 \times \dots \times s_r$ array $\mathcal{A} = (A_{i_1, \dots, i_r})$ (where $r \geq 2$), the *projection* $\psi_{1,2}(\mathcal{A})$ is the $s_1 s_2 \times s_3 \times \dots \times s_r$ array $(B_{i_1, i_3, \dots, i_r})$ given by

$$B_{i_1+s_1 i_2, i_3, \dots, i_r} = A_{i_1, \dots, i_r}, \quad \text{where } 0 \leq i_1 < s_1.$$

The same definition of $\psi_{1,2}$ holds for an array over \mathbb{Z}_3 . For example, $\psi_{1,2}$ maps the three 2×3 arrays over \mathbb{Z}_3 of Example 7 to the three length 6 sequences over \mathbb{Z}_3 of Example 1. For distinct $k, \ell \in \{1, \dots, r\}$, the array $\psi_{k,\ell}(\mathcal{A})$ is defined similarly by replacing the array argument i_ℓ by $i_k + s_k i_\ell$ and removing the array argument i_k . The mapping $\psi_{k,\ell}$ replaces the $s_k \times s_\ell$ “slice” of \mathcal{A} formed from dimensions k and ℓ by the sequence obtained when the elements of the slice are listed column by column.

The following lemma expresses the aperiodic autocorrelation function $C_{\psi_{1,2}(\mathcal{A})}$ as the sum of two terms involving $C_{\mathcal{A}}$ (one or both of which might be trivially zero, according to the values of the arguments). We use this to prove in Theorem 10 that the existence of an r -dimensional Golay array triad implies the existence of an $(r-1)$ -dimensional Golay array triad.

Lemma 9 ([14, Lemma 10]). *Let \mathcal{A} be a 3-phase $s_1 \times \dots \times s_r$ array, where $r \geq 2$. Then*

$$\begin{aligned} C_{\psi_{1,2}(\mathcal{A})}(u_1 + s_1 u_2, u_3, \dots, u_r) \\ = C_{\mathcal{A}}(u_1, \dots, u_r) + C_{\mathcal{A}}(u_1 - s_1, u_2 + 1, u_3, \dots, u_r) \text{ for } 0 \leq u_1 < s_1. \end{aligned}$$

Theorem 10 (Projection mapping). *Suppose that $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a 3-phase $s_1 \times \cdots \times s_r$ Golay array triad, where $r \geq 2$. Then $\{\psi_{1,2}(\mathcal{A}), \psi_{1,2}(\mathcal{B}), \psi_{1,2}(\mathcal{C})\}$ is a 3-phase $s_1 s_2 \times s_3 \times \cdots \times s_r$ Golay array triad.*

Proof. Let u_1, \dots, u_r be integers, where $0 \leq u_1 < s_1$ and $(u_1, \dots, u_r) \neq (0, \dots, 0)$. By Lemma 9,

$$\begin{aligned} & (C_{\psi_{1,2}(\mathcal{A})} + C_{\psi_{1,2}(\mathcal{B})} + C_{\psi_{1,2}(\mathcal{C})})(u_1 + s_1 u_2, u_3, \dots, u_r) \\ &= (C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1, \dots, u_r) + (C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1 - s_1, u_2 + 1, u_3, \dots, u_r) \\ &= 0 \end{aligned}$$

because $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a 3-phase $s_1 \times \cdots \times s_r$ Golay array triad. So $\{\psi_{1,2}(\mathcal{A}), \psi_{1,2}(\mathcal{B}), \psi_{1,2}(\mathcal{C})\}$ is a 3-phase $s_1 s_2 \times s_3 \times \cdots \times s_r$ Golay array triad. \square

The effect of the projection mapping $\psi_{1,2}$ in Theorem 10 is to “join” dimension 1 of a Golay array triad to dimension 2. Likewise, the projection mapping $\psi_{k,\ell}$ for distinct $k, \ell \in \{1, \dots, r\}$ joins dimension k of a Golay array triad to dimension ℓ .

The following corollary arises from repeated application of Theorem 10.

Corollary 11. *Suppose there exists a 3-phase $s_1 \times \cdots \times s_r$ Golay array triad. Then there exists a 3-phase length $\prod_{k=1}^r s_k$ Golay sequence triad.*

Combination of Theorem 4 and Corollary 11 gives the following nonexistence result.

Corollary 12. *There is no 3-phase $s_1 \times \cdots \times s_r$ Golay array triad when $\prod_{k=1}^r s_k \equiv 4 \pmod{6}$.*

We use Theorem 10 to determine the equivalence classes of all 3-phase $s_1 \times \cdots \times s_r$ Golay array triads for which $\prod_{k=1}^r s_k \leq 24$. For example, to determine the equivalence classes of 2×9 Golay array triads, let $\psi_{1,2}$ be the projection mapping from 2×9 arrays to length 18 sequences. Apply the inverse mapping $\psi_{1,2}^{-1}$ to each length 18 Golay sequence triad equivalence class representative (as previously determined, and summarised in Table 1), retaining those 2×9 triads having the Golay property. Collect the resulting Golay array triads into equivalence classes and retain only the representative of each equivalence class. To then determine the equivalence classes of $2 \times 3 \times 3$ Golay triads, let $\psi_{2,3}$ be the projection mapping from $2 \times 3 \times 3$ arrays to 2×9 arrays. Apply the inverse mapping $\psi_{2,3}^{-1}$ to each 2×9 Golay array triad equivalence class representative and proceed similarly. (We can alternatively determine the equivalence classes of $2 \times 3 \times 3$ Golay array triads from the 3×6 Golay array triads, obtained in turn from the length 18 Golay sequence triads.) Likewise, we determine from the length 20 Golay sequence triads that there are no 2×10 and no 4×5 Golay array triads, either of which implies by Theorem 10 that there are also none of size $2 \times 2 \times 5$.

Table 2 shows the resulting counts of equivalence classes of Golay array triads, normalised Golay array triads, and Golay arrays. By Corollary 12, there are no $s_1 \times \cdots \times s_r$ Golay array triads for which $\prod_{k=1}^r s_k \in \{4, 10, 16, 22\}$.

Array size	# equivalence classes								# normalised array triads	# Golay arrays
	size 24	size 48	size 72	size 96	size 144	size 288	size 576	Total		
2×3	1	1						2	72	162
2×4								0	0	0
3×3	1	7		3				11	648	1350
2×6								0	0	0
3×4								0	0	0
2×7						3		3	864	1944
3×5		18		45				63	5184	11664
3×6	4	64	4	147	18	8		245	22464	49788
2×9		18		45	18	18		99	12960	29160
$2 \times 3 \times 3$					2	9	4	15	5184	11664
2×10								0	0	0
4×5								0	0	0
3×7		42		447		9		498	47520	101088
2×12								0	0	0
3×8		6		123				129	12096	27216
4×6								0	0	0

Table 2: Counts of 3-phase $s_1 \times \dots \times s_r$ Golay array triads, where $\prod_{k=1}^r s_k \leq 24$

5 Constructions of Golay array triads

In this section, we present two constructions for Golay array triads, and apply them to a small set of seed Golay sequence and array triads to explain the existence of many of the Golay triads counted in Tables 1 and 2.

The *aperiodic cross-correlation function* of $s_1 \times \dots \times s_r$ arrays $\mathcal{A} = (A_{i_1, \dots, i_r})$ and $\mathcal{B} = (B_{i_1, \dots, i_r})$ is

$$C_{\mathcal{A}, \mathcal{B}}(u_1, \dots, u_r) = \sum_{i_1, \dots, i_r} A_{i_1, \dots, i_r} \overline{B_{i_1+u_1, \dots, i_r+u_r}} \quad \text{for integers } u_1, \dots, u_r$$

(so that $C_{\mathcal{A}, \mathcal{A}}(u_1, \dots, u_r) = C_{\mathcal{A}}(u_1, \dots, u_r)$ for all (u_1, \dots, u_r)). Let $\mathcal{A} = (A_{i_1, \dots, i_r})$, $\mathcal{B} = (B_{i_1, \dots, i_r})$, $\mathcal{C} = (C_{i_1, \dots, i_r})$ be $s_1 \times \dots \times s_r$ arrays. Write $\begin{bmatrix} \mathcal{A} \\ \mathcal{B} \\ \mathcal{C} \end{bmatrix}$ for the $3 \times s_1 \times \dots \times s_r$ array

$\mathcal{D} = (D_{i,i_1,\dots,i_r})$ defined by

$$D_{i,i_1,\dots,i_r} = \begin{cases} A_{i_1,\dots,i_r} & \text{for } i = 0 \\ B_{i_1,\dots,i_r} & \text{for } i = 1 \\ C_{i_1,\dots,i_r} & \text{for } i = 2. \end{cases}$$

It follows directly from the definitions that

$$\left. \begin{aligned} C_{\mathcal{D}}(0, u_1, \dots, u_r) &= (C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1, \dots, u_r), \\ C_{\mathcal{D}}(1, u_1, \dots, u_r) &= (C_{\mathcal{A},\mathcal{B}} + C_{\mathcal{B},\mathcal{C}})(u_1, \dots, u_r), \\ C_{\mathcal{D}}(2, u_1, \dots, u_r) &= C_{\mathcal{A},\mathcal{C}}(u_1, \dots, u_r). \end{aligned} \right\} \quad (5.1)$$

We now present our main construction method for Golay array triads.

Theorem 13 (Increase dimension). *Suppose that $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a 3-phase $s_1 \times \dots \times s_r$ Golay array triad and let $\mathcal{U} = \begin{bmatrix} \mathcal{A} \\ \mathcal{B} \\ \mathcal{C} \end{bmatrix}$, $\mathcal{V} = \begin{bmatrix} \mathcal{A} \\ \omega\mathcal{B} \\ \omega^2\mathcal{C} \end{bmatrix}$, $\mathcal{W} = \begin{bmatrix} \mathcal{A} \\ \omega^2\mathcal{B} \\ \omega\mathcal{C} \end{bmatrix}$. Then $\{\mathcal{U}, \mathcal{V}, \mathcal{W}\}$ is a 3-phase $3 \times s_1 \times \dots \times s_r$ Golay array triad.*

Proof. For all integers u_1, \dots, u_r , by (5.1) we have

$$\begin{aligned} (C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(1, u_1, \dots, u_r) &= (C_{\mathcal{A},\mathcal{B}} + C_{\mathcal{B},\mathcal{C}})(u_1, \dots, u_r) + (C_{\mathcal{A},\omega\mathcal{B}} + C_{\omega\mathcal{B},\omega^2\mathcal{C}})(u_1, \dots, u_r) + \\ &\quad (C_{\mathcal{A},\omega^2\mathcal{B}} + C_{\omega^2\mathcal{B},\omega\mathcal{C}})(u_1, \dots, u_r) \\ &= (1 + \omega^2 + \omega)C_{\mathcal{A},\mathcal{B}}(u_1, \dots, u_r) + (1 + \omega^2 + \omega)C_{\mathcal{B},\mathcal{C}}(u_1, \dots, u_r) \\ &= 0 \end{aligned}$$

and

$$\begin{aligned} (C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(2, u_1, \dots, u_r) &= (C_{\mathcal{A},\mathcal{C}} + C_{\mathcal{A},\omega^2\mathcal{C}} + C_{\mathcal{A},\omega\mathcal{C}})(u_1, \dots, u_r) \\ &= (1 + \omega + \omega^2)C_{\mathcal{A},\mathcal{C}}(u_1, \dots, u_r) \\ &= 0. \end{aligned}$$

Combining these results, we see that the condition

$$(C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(u, u_1, \dots, u_r) = 0 \quad \text{for all integers } u, u_1, \dots, u_r$$

holds for $u \in \{1, 2\}$, and therefore by (4.1) it also holds for $u \in \{-1, -2\}$. It remains to consider the case $u = 0$.

For all integers u_1, \dots, u_r for which $(u_1, \dots, u_r) \neq (0, \dots, 0)$, by (5.1) we have

$$(C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(0, u_1, \dots, u_r)$$

$$\begin{aligned}
&= (C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1, \dots, u_r) + (C_{\mathcal{A}} + C_{\omega\mathcal{B}} + C_{\omega^2\mathcal{C}})(u_1, \dots, u_r) + \\
&\quad (C_{\mathcal{A}} + C_{\omega^2\mathcal{B}} + C_{\omega\mathcal{C}})(u_1, \dots, u_r) \\
&= 3(C_{\mathcal{A}} + C_{\mathcal{B}} + C_{\mathcal{C}})(u_1, \dots, u_r) \\
&= 0
\end{aligned}$$

because $\{\mathcal{A}, \mathcal{B}, \mathcal{C}\}$ is a Golay array triad. Therefore $\{U, V, W\}$ is a 3-phase $3 \times s_1 \times \dots \times s_r$ Golay array triad. \square

Theorem 13 is a simplification and reinterpretation of a construction given by Frank [9, p. 644] for producing a length $3s$ Golay sequence triad over \mathbb{Z}_3 from a length s Golay sequence triad over \mathbb{Z}_3 . We can readily recover Frank's result by applying Theorem 13 to a length s Golay sequence triad, followed by projection of the resulting $3 \times s$ Golay array triad to a length $3s$ Golay sequence triad using Theorem 10. This example illustrates our contention that a higher-dimensional Golay array triad can be considered a more fundamental object than its lower-dimensional projections. Indeed, we now explain many of the entries of Tables 1 and 2 by systematically applying Theorem 13 (which introduces exactly one new dimension) in conjunction with Theorem 10 (which removes exactly one dimension), starting from a small set of seed Golay sequence and array triads.

Before applying Theorem 10 to a set of equivalence class representatives, we replace each representative by its full equivalence class. We then apply the projection mappings $\psi_{k,\ell}$ and $\psi_{\ell,k}$ for all distinct $\{k, \ell\}$ to each element of the class, retaining only the representative of each resulting equivalence class of Golay sequence or array triads. Before applying Theorem 13 to a set of equivalence class representatives, we likewise replace each representative by its full equivalence class, but then also remove the assumption of normalised form and take all $3!$ orderings of the triad sequences (thereby multiplying each class size by a factor of $3^3 \cdot 3!$). We then apply the construction of Theorem 13 to each element of the expanded class, retaining only the representative of each resulting equivalence class of Golay array triads.

Fig. 1 displays the result of applying these two constructions in conjunction. The seed Golay sequence triads are the single equivalence class of length 1 and of length 2 (both of which are trivial), all equivalence classes of length 5, 7 and 8, and one of the 10 equivalence classes of length 6. The seed Golay array triads are nine of the 99 equivalence classes of size 2×9 . Fig. 1 also shows the result of applying Theorem 14 below, which uses length s Golay sequence triads having special cross-correlation properties to construct $2 \times s$ and $3 \times s$ Golay array triads.




Figure 1: Equivalence classes of 3-phase Golay triads whose existence is explained by the constructions of Theorems 13 (increase dimension), 10 (projection mapping), and 14 (cross-correlation). Input seed triads are in red, constructed triads are in blue.

Theorem 14 (Cross-correlation). *Suppose that both $\{\mathcal{A}_1, \mathcal{B}_1, \mathcal{C}_1\}$ and $\{\mathcal{A}_3, \mathcal{B}_3, \mathcal{C}_3\}$ are 3-phase length s Golay sequence triads and that, for all integers u ,*

$$(C_{\mathcal{A}_1, \mathcal{A}_3} + C_{\mathcal{B}_1, \mathcal{B}_3} + C_{\mathcal{C}_1, \mathcal{C}_3})(u) = 0.$$

Then $\left\{ \begin{bmatrix} \mathcal{A}_1 \\ \mathcal{A}_3 \end{bmatrix}, \begin{bmatrix} \mathcal{B}_1 \\ \mathcal{B}_3 \end{bmatrix}, \begin{bmatrix} \mathcal{C}_1 \\ \mathcal{C}_3 \end{bmatrix} \right\}$ is a 3-phase $2 \times s$ Golay array triad.

Suppose that $\{\mathcal{A}_2, \mathcal{B}_2, \mathcal{C}_2\}$ is also a 3-phase length s Golay sequence triad and that, for all integers u ,

$$(C_{\mathcal{A}_1, \mathcal{A}_2} + C_{\mathcal{A}_2, \mathcal{A}_3} + C_{\mathcal{B}_1, \mathcal{B}_2} + C_{\mathcal{B}_2, \mathcal{B}_3} + C_{\mathcal{C}_1, \mathcal{C}_2} + C_{\mathcal{C}_2, \mathcal{C}_3})(u) = 0.$$

Then $\left\{ \begin{bmatrix} \mathcal{A}_1 \\ \mathcal{A}_2 \\ \mathcal{A}_3 \end{bmatrix}, \begin{bmatrix} \mathcal{B}_1 \\ \mathcal{B}_2 \\ \mathcal{B}_3 \end{bmatrix}, \begin{bmatrix} \mathcal{C}_1 \\ \mathcal{C}_2 \\ \mathcal{C}_3 \end{bmatrix} \right\}$ is a 3-phase $3 \times s$ Golay array triad.

Proof. We give the proof for the $3 \times s$ array triad; the proof for the $2 \times s$ array triad is similar. Let $\mathcal{U} = \begin{bmatrix} \mathcal{A}_1 \\ \mathcal{A}_2 \\ \mathcal{A}_3 \end{bmatrix}$, $\mathcal{V} = \begin{bmatrix} \mathcal{B}_1 \\ \mathcal{B}_2 \\ \mathcal{B}_3 \end{bmatrix}$, $\mathcal{W} = \begin{bmatrix} \mathcal{C}_1 \\ \mathcal{C}_2 \\ \mathcal{C}_3 \end{bmatrix}$. For all integers $u \neq 0$, by (5.1) we have

$$\begin{aligned} & (C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(0, u) \\ &= (C_{\mathcal{A}_1} + C_{\mathcal{A}_2} + C_{\mathcal{A}_3})(u) + (C_{\mathcal{B}_1} + C_{\mathcal{B}_2} + C_{\mathcal{B}_3})(u) + (C_{\mathcal{C}_1} + C_{\mathcal{C}_2} + C_{\mathcal{C}_3})(u) = 0 \end{aligned}$$

because each of $\{\mathcal{A}_1, \mathcal{B}_1, \mathcal{C}_1\}$, $\{\mathcal{A}_2, \mathcal{B}_2, \mathcal{C}_2\}$, $\{\mathcal{A}_3, \mathcal{B}_3, \mathcal{C}_3\}$ is a Golay sequence triad. Furthermore, for all integers u we have

$$\begin{aligned} & (C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(1, u) \\ &= (C_{\mathcal{A}_1, \mathcal{A}_2} + C_{\mathcal{A}_2, \mathcal{A}_3})(u) + (C_{\mathcal{B}_1, \mathcal{B}_2} + C_{\mathcal{B}_2, \mathcal{B}_3})(u) + (C_{\mathcal{C}_1, \mathcal{C}_2} + C_{\mathcal{C}_2, \mathcal{C}_3})(u) = 0 \end{aligned}$$

by assumption, and

$$(C_{\mathcal{U}} + C_{\mathcal{V}} + C_{\mathcal{W}})(2, u) = C_{\mathcal{A}_1, \mathcal{A}_3}(u) + C_{\mathcal{B}_1, \mathcal{B}_3}(u) + C_{\mathcal{C}_1, \mathcal{C}_3}(u) = 0$$

by assumption. It follows from (4.1) that $\{\mathcal{U}, \mathcal{V}, \mathcal{W}\}$ is a 3-phase $3 \times s$ Golay array triad. \square

To construct 3×7 and 2×7 Golay array triads using Theorem 14, note that the three (inequivalent) 3-phase length 7 Golay sequence triads

$$\begin{aligned} \mathcal{T}_1 &= \{\mathcal{U}_1, \mathcal{V}_1, \mathcal{W}_1\} = \left\{ \left[\begin{array}{ccccccc} 1 & 1 & \omega & 1 & \omega^2 & \omega^2 & 1 \end{array} \right], \left[\begin{array}{ccccccc} 1 & 1 & 1 & \omega^2 & 1 & \omega & \omega \end{array} \right], \left[\begin{array}{ccccccc} 1 & 1 & \omega^2 & 1 & \omega & 1 & \omega^2 \end{array} \right] \right\}, \\ \mathcal{T}_2 &= \{\mathcal{U}_2, \mathcal{V}_2, \mathcal{W}_2\} = \left\{ \left[\begin{array}{ccccccc} 1 & \omega & \omega & \omega & 1 & \omega & \omega^2 \end{array} \right], \left[\begin{array}{ccccccc} 1 & \omega & 1 & \omega^2 & \omega & 1 & 1 \end{array} \right], \left[\begin{array}{ccccccc} 1 & \omega & \omega^2 & \omega^2 & \omega^2 & \omega^2 & \omega \end{array} \right] \right\}, \\ \mathcal{T}_3 &= \{\mathcal{U}_3, \mathcal{V}_3, \mathcal{W}_3\} = \left\{ \left[\begin{array}{ccccccc} 1 & \omega^2 & \omega & \omega & \omega & 1 & \omega \end{array} \right], \left[\begin{array}{ccccccc} 1 & \omega^2 & 1 & \omega & \omega^2 & \omega^2 & \omega^2 \end{array} \right], \left[\begin{array}{ccccccc} 1 & \omega^2 & \omega^2 & 1 & 1 & \omega & 1 \end{array} \right] \right\} \end{aligned}$$

satisfy, for all integers u ,

$$\begin{aligned} (C_{\mathcal{U}_1, \mathcal{U}_2} + C_{\mathcal{V}_1, \mathcal{V}_2} + C_{\mathcal{W}_1, \mathcal{W}_2})(u) &= 0, \\ (C_{\mathcal{U}_2, \mathcal{U}_3} + C_{\mathcal{V}_2, \mathcal{V}_3} + C_{\mathcal{W}_2, \mathcal{W}_3})(u) &= 0, \\ (C_{\mathcal{U}_3, \mathcal{U}_1} + C_{\mathcal{V}_3, \mathcal{V}_1} + C_{\mathcal{W}_3, \mathcal{W}_1})(u) &= 0. \end{aligned}$$

We may therefore apply Theorem 14 to construct three 3-phase 2×7 Golay array triads by taking $(\{\mathcal{A}_1, \mathcal{B}_1, \mathcal{C}_1\}, \{\mathcal{A}_3, \mathcal{B}_3, \mathcal{C}_3\})$ to be $(\mathcal{T}_1, \mathcal{T}_2)$ or $(\mathcal{T}_2, \mathcal{T}_3)$ or $(\mathcal{T}_3, \mathcal{T}_1)$, and these three constructed triads are inequivalent. Using the identity

$$C_{\mathcal{B}, \mathcal{A}}(u) = \overline{C_{\mathcal{A}, \mathcal{B}}(-u)} \quad \text{for all } u$$

for complex-valued sequences \mathcal{A}, \mathcal{B} of equal length, we may also apply Theorem 14 to construct nine 3-phase 3×7 Golay array triads by taking $(\{\mathcal{A}_1, \mathcal{B}_1, \mathcal{C}_1\}, \{\mathcal{A}_2, \mathcal{B}_2, \mathcal{C}_2\}, \{\mathcal{A}_3, \mathcal{B}_3, \mathcal{C}_3\})$ to be $(\omega^e \mathcal{T}_1, \mathcal{T}_2, \mathcal{T}_3)$ or $(\mathcal{T}_2, \mathcal{T}_3, \omega^e \mathcal{T}_1)$ or $(\mathcal{T}_3, \omega^e \mathcal{T}_1, \mathcal{T}_2)$ for each $e \in \mathbb{Z}_3$, and these nine constructed triads are inequivalent.

We see from Fig. 1 that, starting from a small set of seed Golay sequence and array triads, we can use Theorems 10, 13, and 14 to explain the existence of all 2×3 , 3×3 , 2×7 , 3×5 , 3×6 , $2 \times 3 \times 3$, 3×7 , and 3×8 Golay array triads, and all but nine of the 99 equivalence classes of 2×9 Golay array triads. We can also explain a large proportion of the equivalence classes of length 3, 6, 9, 14, 15, 18, 21, and 24 Golay sequence triads, including all but nine of the 3945 equivalence classes of length 21 and all but five of the 1025 equivalence classes of length 24. The counts of equivalence classes recorded in Tables 1 and 2 that are not explained by these constructions are summarised in Table 3. The representative of some of these unexplained equivalence classes is listed in the Appendix.

6 Open Questions

In this section, we pose some open questions motivated by our results. The counts shown in Table 1, and the nonexistence result of Theorem 4, suggest a natural question:

Q1. Does there exist a 3-phase length s Golay sequence triad if and only if $s \not\equiv 4 \pmod{6}$?

The counts shown in Table 2 suggest a possible generalisation of Theorem 4 to each of the dimensions of a Golay array triad:

Q2. Does the existence of a 3-phase $s_1 \times \dots \times s_r$ Golay array triad imply that $s_k \not\equiv 4 \pmod{6}$ for each k ?

The repetition of some of the counts shown in Table 2 suggests a possible connection:

Q3. Is there a convincing explanation for the total count of normalised array triads and of Golay arrays being identical for 3-phase Golay triads of sizes 3×5 and $2 \times 3 \times 3$?

The unexplained equivalence classes counted in Table 3 prompt the question:

Q4. Can new constructions be found to account for the equivalence classes of Golay sequence and array triads marked as “some” or “none” in Table 3?

Sequence or array size	Total # equivalence classes	# unexplained equivalence classes					Total	none/some/all explained, or seeds
		size 1	size 24	size 48	size 288			
2	1	1					1	trivial seed
5	3		3				3	seeds
7	17		8	9			17	seeds
8	4		1	3			4	seeds
11	64		14	50			64	none
12	7			7			7	none
13	64		16	48			64	none
17	25		10	15			25	none
19	17		6	11			17	none
20	10			10			10	none
23	2		2				2	none
3	2						0	all
6	10		1				1	some (*)
9	105		10	26			36	some
14	91		23	32			55	some
15	519		15	36			51	some
18	1485		28	5			33	some
21	3945		4	5			9	some
24	1025		4	1			5	some
2×3	2						0	all
3×3	11						0	all
2×7	3						0	all
3×5	63						0	all
3×6	245						0	all
2×9	99				9		9	some (*)
$2 \times 3 \times 3$	15						0	all
3×7	498						0	all
3×8	129						0	all

Table 3: Counts of 3-phase Golay sequence and array triads whose existence is not explained by the constructions of Theorems 10, 13, 14. (*) indicates that the unexplained equivalence class(es) are used as seeds for other sizes.

Appendix: Unexplained Golay sequence and array triads

In this Appendix, we give the representative of those equivalence classes of Golay triads that are not explained by the constructions of Section 5, for lengths 6, 21, and 24, and for size 2×9 .

The single unexplained equivalence class of length 6 Golay sequence triads over \mathbb{Z}_3 has representative

$$\{ [0\ 0\ 0\ 1\ 1\ 0], [0\ 2\ 0\ 2\ 2\ 1], [0\ 1\ 2\ 2\ 0\ 2] \}.$$

The nine unexplained equivalence classes of length 21 Golay sequence triads over \mathbb{Z}_3 have representatives

$$\begin{aligned} & \{ [000012012221021212210], [011200122121112120221], [000021101021102000012] \}, \\ & \{ [000012210120010100210], [000012210001201212121], [000012210222122021002] \}, \\ & \{ [000122212210000110010], [000121002010112021221], [000120020110020202102] \}, \\ & \{ [001101021212120101100], [001100020011211010221], [002200010022122020112] \}, \\ & \{ [001102202020212201100], [001122210020201211211], [001001001010201110022] \}, \\ & \{ [001120012210202100210], [000022220000201012121], [002122120100120020002] \}, \\ & \{ [011110012220121122210], [000212112010201002101], [000121022022022220102] \}, \\ & \{ [011121121201002221210], [000220112222102210201], [001102000001201210102] \}, \\ & \{ [011122022122111021210], [000111102212211212101], [001100120012120100202] \}. \end{aligned}$$

The five unexplained equivalence classes of length 24 Golay sequence triads over \mathbb{Z}_3 have representatives

$$\begin{aligned} & \{ [000010100112012002112100], [001222010101111102100221], [001100220110200202021212] \}, \\ & \{ [000011110101200220122210], [000212010020122002101021], [000112110212011110010102] \}, \\ & \{ [000100011012212110201210], [000120002200201101022021], [000222201121220201021102] \}, \\ & \{ [012022202000212221011210], [000121101221020002022201], [000020110012100121001102] \}, \\ & \{ [012122010210012010221210], [000200022112002122220201], [000100011221001211110102] \}. \end{aligned}$$

The nine unexplained equivalence classes of 2×9 Golay array triads over \mathbb{Z}_3 have representatives

$$\begin{aligned} & \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 1 & 2 \\ 2 & 1 & 0 & 2 & 0 & 1 & 2 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 & 0 & 0 & 1 & 1 & 2 & 1 & 2 \\ 1 & 2 & 2 & 1 & 1 & 0 & 2 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 & 2 & 0 & 0 & 1 & 0 & 1 \\ 2 & 2 & 1 & 1 & 0 & 1 & 0 & 0 & 2 \end{bmatrix} \right\}, \\ & \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 2 & 2 & 2 \\ 2 & 1 & 0 & 2 & 0 & 2 & 1 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 & 2 & 1 & 1 & 2 & 1 & 2 \\ 1 & 1 & 0 & 0 & 1 & 0 & 2 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 & 0 & 2 & 1 & 1 & 0 & 0 \\ 1 & 1 & 2 & 1 & 2 & 0 & 1 & 1 & 2 \end{bmatrix} \right\}, \end{aligned}$$

$$\begin{aligned}
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 2 \\ 2 & 0 & 1 & 2 & 1 & 1 & 0 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 0 & 2 & 1 & 1 & 1 & 2 \\ 1 & 2 & 2 & 1 & 2 & 0 & 2 & 1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 2 & 0 & 0 & 1 & 1 & 1 \\ 2 & 0 & 2 & 1 & 0 & 1 & 1 & 0 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 2 & 2 & 2 \\ 0 & 1 & 2 & 0 & 0 & 1 & 2 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 2 & 0 & 0 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 & 2 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 1 & 2 & 1 & 0 & 0 & 1 & 1 \\ 0 & 2 & 0 & 2 & 0 & 1 & 0 & 0 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 \\ 1 & 2 & 0 & 1 & 0 & 1 & 0 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 0 & 0 & 2 & 2 & 2 & 0 \\ 0 & 1 & 1 & 0 & 2 & 0 & 2 & 1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 2 & 0 & 1 & 2 & 2 & 2 \\ 1 & 2 & 1 & 0 & 2 & 1 & 1 & 0 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 0 & 1 & 2 & 0 & 1 & 1 \\ 1 & 0 & 2 & 1 & 1 & 1 & 0 & 2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 0 & 2 & 2 & 0 & 1 & 0 \\ 2 & 1 & 2 & 2 & 2 & 0 & 2 & 1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 2 & 0 & 2 & 1 & 2 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2 \\ 0 & 2 & 1 & 1 & 2 & 1 & 2 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 2 & 1 & 0 & 2 & 2 & 0 \\ 0 & 2 & 0 & 2 & 0 & 1 & 1 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 0 & 2 & 0 & 1 & 1 & 1 \\ 0 & 2 & 2 & 0 & 1 & 1 & 0 & 1 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 1 & 0 & 2 & 0 \\ 1 & 2 & 0 & 2 & 2 & 1 & 2 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 2 & 1 & 1 & 2 & 1 & 1 \\ 1 & 2 & 2 & 0 & 0 & 1 & 1 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 0 & 2 & 1 & 1 & 0 & 2 \\ 1 & 2 & 1 & 1 & 1 & 1 & 0 & 1 & 2 \end{bmatrix} \right\}, \\
& \left\{ \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 2 & 1 & 0 & 1 \\ 0 & 1 & 2 & 1 & 1 & 1 & 2 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 2 & 2 & 1 & 2 & 0 & 2 & 2 \\ 0 & 1 & 1 & 2 & 2 & 1 & 1 & 2 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 & 0 & 2 & 2 & 2 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 2 \end{bmatrix} \right\}.
\end{aligned}$$

References

- [1] A.A. Avis. 3-phase Golay triads. Master's thesis, Simon Fraser University, 2008. Available online: <http://summit.sfu.ca/item/11482>.
- [2] J.P. Bell, J. Jedwab, M. Khatirinejad, and K.-U. Schmidt. Three-phase Barker arrays. *J. Comb. Des.*, **23**:45–59, 2015.
- [3] C.V. Chong, R. Venkataramani, and V. Tarokh. A new construction of 16-QAM Golay complementary sequences. *IEEE Trans. Inform. Theory*, **49**:2953–2959, 2003.
- [4] R. Craigen, W. Holzmann, and H. Kharaghani. Complex Golay sequences: structure and applications. *Discrete Math.*, **252**:73–89, 2002.
- [5] J.A. Davis and J. Jedwab. Peak-to-mean power control in OFDM, Golay complementary sequences, and Reed-Muller codes. *IEEE Trans. Inform. Theory*, **45**:2397–2417, 1999.
- [6] M. Dymond. *Barker arrays: existence, generalization and alternatives*. PhD thesis, University of London, 1992.
- [7] F. Fiedler, J. Jedwab, and M.G. Parker. A multi-dimensional approach to the construction and enumeration of Golay complementary sequences. *J. Combin. Theory (A)*, **115**:753–776, 2008.

- [8] F. Fiedler, J. Jedwab, and A. Wiebe. A new source of seed pairs for Golay sequences of length 2^m . *J. Combin. Theory (A)*, **117**:589–597, 2010.
- [9] R.L. Frank. Polyphase complementary codes. *IEEE Trans. Inform. Theory*, **IT-26**:641–647, 1980.
- [10] R.G. Gibson and J. Jedwab. Quaternary Golay sequence pairs I: Even length. *Des. Codes Cryptogr.*, **59**:131–146, 2011.
- [11] M.J.E. Golay. Static multislit spectrometry and its application to the panoramic display of infrared spectra. *J. Opt. Soc. Amer.*, **41**:468–472, 1951.
- [12] M.J.E. Golay. Complementary series. *IRE Trans. Inform. Theory*, **IT-7**:82–87, 1961.
- [13] J. Jedwab. What can be used instead of a Barker sequence? *Contemp. Math.*, **461**:153–178, 2008.
- [14] J. Jedwab and M.G. Parker. Golay complementary array pairs. *Des. Codes Cryptogr.*, **44**:209–216, 2007.
- [15] M. Nazarathy, S.A. Newton, R.P. Giffard, D.S. Moberly, F. Sischka, W.R. Trutna, Jr., and S. Foster. Real-time long range complementary correlation optical time domain reflectometer. *IEEE J. Lightwave Technology*, **7**:24–38, 1989.
- [16] A. Nowicki, W. Secomski, J. Litniewski, I. Trots, and P.A. Lewin. On the application of signal compression using Golay’s codes sequences in ultrasonic diagnostic. *Arch. Acoustics*, **28**:313–324, 2003.
- [17] N. Ohyama, T. Honda, and J. Tsujiuchi. An advanced coded imaging without side lobes. *Optics Comm.*, **27**:339–344, 1978.
- [18] K.G. Paterson. Generalized Reed-Muller codes and power control in OFDM modulation. *IEEE Trans. Inform. Theory*, **46**:104–120, 2000.
- [19] C. Rößing and V. Tarokh. A construction of OFDM 16-QAM sequences having low peak power. *IEEE Trans. Inform. Theory*, **47**:2091–2094, 2001.
- [20] K.-U. Schmidt. Complementary sets, generalized Reed-Muller codes, and power control for OFDM. *IEEE Trans. Inform. Theory*, **53**:808–814, 2007.
- [21] R.J. Turyn. Hadamard matrices, Baumert-Hall units, four-symbol sequences, pulse compression, and surface wave encodings. *J. Combin. Theory (A)*, **16**:313–333, 1974.